
**U.S. Department
of Commerce**
Seattle, Washington

**Volume 101
Number 4
October 2003**

Fishery Bulletin

Contents

Articles

- 721–731 **Baum, Julia K., Jessica J. Meeuwig, and
Amanda C. J. Vincent**
Bycatch of lined seahorses (*Hippocampus erectus*) in a
Gulf of Mexico shrimp trawl fishery
- 732–736 **Bjorndal, Karen A., Alan B. Bolten, and Helen R. Martins**
Estimates of survival probabilities for oceanic-stage loggerhead
sea turtles (*Caretta caretta*) in the North Atlantic
- 737–744 **Chen, Yong, Margaret Hunter, Robert Vadas, and
Brian Beal**
Developing a growth-transition matrix for the stock assessment of
the green sea urchin (*Strongylocentrotus droebachiensis*) off Maine
- 745–757 **de Lestang, Simon, Norman G. Hall, and Ian C. Potter**
Reproductive biology of the blue swimmer crab (*Portunus pelagicus*,
Decapoda: Portunidae) in five bodies of water on the west coast of
Australia
- 758–768 **Dew, Jodi R., Jim Berkson, Eric M. Hallerman, and
Standish K. Allen Jr.**
A model for assessing the likelihood of self-sustaining populations
resulting from commercial production of triploid Suminoe oysters
(*Crassostrea ariakensis*) in Chesapeake Bay
- 769–777 **Díaz-Jaimes, Pindaro, and Manuel Uribe-Alcocer**
Allozyme and RAPD variation in the eastern Pacific yellowfin tuna
(*Thunnus albacares*)
- 778–789 **Govoni, John Jeffrey, Elisabeth H. Laban, and
Jonathan A. Hare**
The early life history of swordfish (*Xiphias gladius*) in the western
North Atlantic
- 790–799 **Heales, Donald S., David T. Brewer, You-Gan Wang, and
Peter N. Jones**
Does the size of subsamples take from multispecies trawl catches
affect estimates of catch composition and abundance?

The conclusions and opinions expressed in *Fishery Bulletin* are solely those of the authors and do not represent the official position of the National Marine Fisheries Service (NOAA) or any other agency or institution.

The National Marine Fisheries Service (NMFS) does not approve, recommend, or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

- 800–808** **Laidig, Thomas E., Donald E. Pearson, and Lorraine L. Sinclair**
Age and growth of blue rockfish (*Sebastes mystinus*) from central and northern California
- 809–821** **Marin E., Baumar J., Antonio Quintero, Dany Bussière, and Julian J. Dodson**
Reproduction and recruitment of white mullet (*Mugil curema*) to a tropical lagoon (Margarita Island, Venezuela) as revealed by otolith microstructure
- 822–834** **McDonough, Christopher J., William A. Roumillat, and Charles A. Wenner**
Fecundity and spawning season of striped mullet (*Mugil cephalus* L.) in South Carolina estuaries
- 835–850** **Nelson, Peter A.**
Marine fish assemblages associated with fish aggregating devices (FADs): effects of fish removal, FAD size, fouling communities, and prior recruits
- 851–859** **Pajuelo, José G., José M. Lorenzo, and Muriel Gregoire**
Age and growth of the bastard grunt (*Pomadasys incisus*) Haemulidae inhabiting the Canarian archipelago, Northwest Africa
- 860–873** **Punt, André E.**
Evaluating the efficacy of managing West Coast groundfish resources through simulations
- 874–888** **Purves, Martin G., David J. Agnew, Guillermo Moreno, Tim Daw, Cynthia Yau, and Graham Pilling**
Distribution, demography, and discard mortality of crabs caught as bycatch in an experimental pot fishery for toothfish (*Dissostichus eleginoides*) in the South Atlantic
- 889–899** **Stabenau, Erich K., and Kimberly R. N. Vietti**
The physiological effects of multiple forced submergences in loggerhead sea turtles (*Caretta caretta*)
- 900–909** **Stephenson, Peter C., and Norm G. Hall**
Quantitative determination of the timing of otolith ring formation from marginal increments in four marine teleost species from northwestern Australia

Notes

- 910–914** **Chan, Ricky W. K., Patricia I. Dixon, Julian G. Pepperell, and Dennis D. Reid**
Application of DNA-based techniques for the identification of whaler sharks (*Carcharhinus* spp.) caught in protective beach meshing and by recreational fisheries off the coast of New South Wales
- 915–922** **Ebert, Thomas A., and John R. Southon**
Red sea urchins (*Strongylocentrotus franciscanus*) can live over 100 years: confirmation with A-bomb ¹⁴carbon
- 923–932** **Fulling, Gregory L., Keith D Mullin, and Carrie W. Hubard**
Abundance and distribution of cetaceans in outer continental shelf waters of the U.S. Gulf of Mexico
- 933–938** **Hata, David, and Jim Berkson**
Abundance of horseshoe crabs (*Limulus polyphemus*) in the Delaware Bay area
- 939–948** **Kerstetter, David W., Brian E. Luckhurst, Eric D. Prince, and John E. Graves**
Use of pop-up satellite archival tags to demonstrate survival of blue marlin (*Makaira nigricans*) released from pelagic longline gear
- 949–950** *2003 reviewers*
- 951–960** *2003 index*
- 961** *Subscription form*